

For adults and children 6 months of age and older with dystrophic or junctional epidermolysis bullosa (EB)

Are you considering FILSUEZ?

Here's what you should know



Henry, a real FILSUEZ patient

Important Safety Information

What is FILSUEZ (birch triterpenes) topical gel for?


FILSUEZ is a prescription medicine used on the skin to treat wounds that may happen with dystrophic and junctional epidermolysis bullosa (EB) in adults and children 6 months of age and older.

It is not known if FILSUEZ is safe and effective in children younger than 6 months of age.

Please see additional **Important Safety Information** throughout.
Please see full **Product Information** and **Patient Information**.

FILSUVÉZ was shown to improve wound healing

In a clinical study of 223 people with EB (195 with dystrophic EB and 26 with junctional EB)



Target wounds closed for **41.3%** of people using FILSUVÉZ within **45 days** compared with **28.9%** of those using placebo gel.



You may experience wound healing and closure **in as little as 45 days** after starting FILSUVÉZ.

Important Safety Information (cont'd)

What is the most important safety information I should know?

FILSUVÉZ may cause a serious side effect of allergic reactions.

Allergic reactions and skin reactions to FILSUVÉZ may include the following symptoms: red itchy bumps (hives), skin rash, redness or itching. If you get any of these symptoms, stop using FILSUVÉZ right away and call your healthcare provider.

Please see additional Important Safety Information throughout.

Please see full Product Information and Patient Information.

FILSUVEZ was proven to be safe and well tolerated



The most common side effect of FILSUVEZ is application site reactions, such as pain and itchy skin

If the wounds you are treating with FILSUVEZ become infected, stop treatment and contact your doctor.

- Possible signs of infection include, but are not limited to, the wound becoming red, warm, swollen, painful, and/or draining yellow or greenish fluids (pus)

Important Safety Information (cont'd)

What are the possible side effects of FILSUVEZ?

The most common side effect of FILSUVEZ is application site reactions, such as pain and itchy skin.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088

How should I take FILSUVEZ?

See the detailed “Instructions for Use” that comes with FILSUVEZ for information on how to apply FILSUVEZ.

Use FILSUVEZ exactly as your healthcare provider tells you to use it.

The tube of FILSUVEZ is for one-time use only. After the tube has been opened, apply the gel right away. Throw away any remaining gel and the tube after use.

Please see additional Important Safety Information throughout.

Please see full Product Information and Patient Information.

FILSUVÉZ is a topical gel that can be easily added to your treatment routine

Applied from the comfort of your own home to **every wound**, at **every wound dressing change**.



[**Click here to see how FILSUVÉZ supported Henry, who lives with dystrophic EB**](#)



Important Safety Information (cont'd)

How should I take FILSUVÉZ? (cont'd)

Do not use around or get FILSUVÉZ in the eyes, or mucous membrane areas, examples are mouth, vagina or anus.

If you get FILSUVÉZ in your eyes or mucous membrane area, rinse with clean water right away. Contact your healthcare provider if you have any discomfort.

If the wounds you are treating with FILSUVÉZ become infected, **stop treatment** and contact your healthcare provider. Signs or symptoms of infection may include the wound becoming red, warm, swollen, painful or drains yellow or greenish fluid (pus).

Please see additional [Important Safety Information](#) throughout.

Please see full [Product Information](#) and [Patient Information](#).



©2025 Chiesi USA. All rights reserved.

Filsuvez® is a registered trademark owned by the Chiesi Group.

PP-RA-00698 V1.0